

A new method of measuring the cluster peculiar velocity power spectrum

Pengjie Zhang^{1,2,*}, Hume A. Feldman^{3,†}, Roman Juszkiewicz^{4,5,‡} & Albert Stebbins^{2,*}

¹*Shanghai Observatory, Chinese Academy of Science, Shanghai, 200030 China*

²*NASA/Fermilab Astrophysics Center, Fermi National Accelerator Laboratory, Batavia, IL 60510–0500*

³*Dept. of Physics & Astronomy, Univ. of Kansas, Lawrence, KS 66045, USA*

⁴*Institute of Astronomy, Zielona Góra University, 65–516 Zielona Góra, Poland*

⁵*Copernicus Astronomical Center, 00–716 Warsaw, Poland*

*emails: *pjzhang@shao.ac.cn; †feldman@ku.edu; ‡roman@camk.edu.pl; *stebbins@fnal.gov*

25 June 2008

ABSTRACT

We propose to use spatial correlations of the kinetic Sunyaev–Zeldovich (KSZ) flux as an estimator of the peculiar velocity power spectrum. In contrast with conventional techniques, our new method does not require measurements of the thermal SZ signal or the X–ray temperature. Moreover, this method has the special advantage that the expected systematic errors are always sub–dominant to statistical errors on all scales and redshifts of interest. We show that future large sky coverage KSZ surveys may allow a peculiar velocity power spectrum estimates of an accuracy reaching $\sim 10\%$.

Subject headings: cosmology: large scale structure; theory–cosmic microwave background; cosmology: observations; cosmology: theory; dark matter; distance scale

1 INTRODUCTION

Peculiar velocity–distance galaxy surveys have provided us with maps of line-of-sight peculiar velocity fields, $v(\mathbf{x}) = \mathbf{V} \cdot \mathbf{x}/x$, where $\mathbf{V}$ is the peculiar velocity at position $\mathbf{x}$. Such surveys were successfully used in the past to constrain power spectra of initial density fluctuations and to estimate the cosmological density parameter, Ω_m (Vittorio, Juszkiewicz & Davis 1986; Groth, Juszkiewicz & Ostriker 1989; da Costa et al. 1998; Feldman, et al. 2003a; Feldman, et al. 2003b; Sarkar, Feldman & Watkins 2007; Watkins & Feldman 2007; Feldman & Watkins 2008). For an overview of the subject, see (Strauss & Willick 1995) and (Courteau & Willick 2000). However, all these velocity–distance surveys are limited by errors in the range of 15 to 20% of the estimated distance. Measurements based on type Ia supernovae have slightly lower error per source, due to smaller dispersions in type Ia supernovae intrinsic luminosities (see, e.g. (Hui & Greene 2006; Gordon, Land & Slosar 2007) and references therein for theory aspects and (Bonvin, Durrer & Kunz, 2006; Haugboelle et al. 2007; 1; Wang 2007; Watkins & Feldman 2007) and references therein for recent measurements). In contrast, for cluster line-of-sight peculiar velocities, v , derived from the kinetic Sunyaev Zeldovich (KSZ) effect, the errors grow much less rapidly with distance. KSZ cluster surveys may open new possibilities for studying large–scale flows (Haehnelt & Tegmark 1996; Kashlinsky & Atrio-Barandela 2000; Aghanim, Gorski & Puget 2001; Atrio-Barandela, Kashlinsky & Mucket, 2004; Holder

2004) and cosmology (Bhattacharya & Kosowsky, 2007; Fos-alba & Doré 2007). The currently established method of recovering v from the SZ data requires extra measurements of the cluster temperature and its Thomson optical depth, τ . Other complications include the relativistic thermal Sunyaev Zeldovich (TSZ) effect and the KSZ signal generated by internal motions, which is added to the signal from cluster’s bulk velocity, resulting in a limit in accuracy in determining v , of the order of ~ 200 km/s (Knox, Holder & Church 2004; Aghanim, Hansen & Lagache, 2005; Diaferio et al. 2005).

In this paper we propose to estimate the spatial two–point correlation function of the KSZ flux from the data and then infer the velocity power spectrum. Contaminations to the cluster KSZ flux have various clustering properties that can be applied to disentangle the signal from its contaminants (see also (Zhang et al. 2004)). We will show that at $z \gtrsim 0.3$, the systematics become sub–dominant and the statistical errors, at $\sim 10\%$ level for South Pole telescope (SPT¹) (Ruhl et al. 2004), dominate. Throughout this *letter*, we adopt the cosmology with $\Omega_m = 0.3$, $\Omega_\Lambda = 1 - \Omega_m$ and a normalization parameter $\sigma_8 = 0.9$.

2 THE FLUX POWER SPECTRUM

The KSZ cluster surveys directly measure the sum of the cluster KSZ flux S_{KSZ} and various contaminants, such as intracluster gas internal flow, radio and IR point sources, primary cosmic microwave background (CMB), cosmic infrared background (CIB), etc. The signal is (Sunyaev & Zeldovich 1980)

¹ <http://spt.uchicago.edu/>, specifically, see the SPT White Paper at this site.

$$S_{\text{KSZ}} = \frac{\partial B_\nu(T)}{\partial T} \Delta T_{\text{KSZ}}, \quad (1)$$

where $B_\nu(T)$ is the Planck function. At $\nu \sim 217\text{GHz}$, $\partial B/\partial T = 540 \text{ Jy sr}^{-1} \mu\text{K}^{-1}$. The KSZ temperature fluctuation can be expressed as

$$\Delta T_{\text{KSZ}} = 2.7\text{K} \langle \tau \rangle (v/c) = 9\mu\text{K} v_{100} \langle \tau \rangle / 0.01, \quad (2)$$

where $\langle \tau \rangle$ is the optical depth averaged over the solid angle ω , while $v_{100} \equiv v/(100\text{km/s})$. We will also use $S_{100} \equiv S_{\text{KSZ}}(v)/S_{\text{KSZ}}(100\text{km/s})$. These quantities are, respectively, the “normalized” peculiar velocity and KSZ flux, scaled to $v = 100\text{km/s}$. The value $\langle \tau \rangle = 0.01$ is typical for cluster models considered in numerical simulations (see e.g. (Yoshida, Sheth & Diaferio 2001) and references therein). Since at $\nu \sim 217\text{GHz}$, the non-relativistic part of the TSZ effect, which is one of the major contaminants of the KSZ, vanishes (Zel’dovich & Sunyaev 1969), we focus on this frequency. Optical follow-up of KSZ surveys such as dark energy survey² will measure cluster redshift z with uncertainty $\lesssim 0.005$.³ The z information allows the measurement of the 3D auto correlation of measured cluster KSZ flux

$$\xi_S(r) \equiv \langle \Delta S_i \Delta S_j \rangle = \langle S_{\text{KSZ},i} S_{\text{KSZ},j} \rangle + \dots, \quad (3)$$

where $\Delta S_i = S_i - \bar{I}\omega_i$, S_i and ω_i are the total flux and the solid angle subtended by the i -th cluster; $r = |\mathbf{x}_i - \mathbf{x}_j|$ is the comoving separation of a pair of clusters, and $\bar{I} = \sum S_i / \sum \omega_i$ is the mean flux intensity. The total flux is a sum of the KSZ signal, $S_{\text{KSZ},i}$, and various noise terms, which we study below.

The correlation $\langle S_{\text{KSZ},i} S_{\text{KSZ},j} \rangle$ is cluster number density weighted and thus

$$\langle S_{\text{KSZ},i} S_{\text{KSZ},j} \rangle \propto \langle (1 + \delta_C(M_i)) v_i (1 + \delta_C(M_j)) v_j \rangle, \quad (4)$$

where $\delta_C(M) = b_n(M)\delta$ is the cluster number overdensity, while δ is the matter overdensity. Here we use a simple biasing model, where the parameter b_n is determined by the cluster mass M (Mo & White 1996) (see also refs. (Kashlinsky 1987; Kashlinsky 1991)). The quantity above can be written as

$$\langle (1 + \delta_i) v_i (1 + \delta_j) v_j \rangle = \langle v_i v_j \rangle + \langle \delta_i \delta_j v_i v_j \rangle, \quad (5)$$

where the third order moments cancel because of statistical isotropy and homogeneity which reflects the symmetry expected in the large scale pattern of the cluster distribution: $\langle v_i v_j \delta_i \rangle = -\langle v_i v_j \delta_j \rangle$. The fourth moment can always be written as a sum of products of second moments of all possible pairs plus the irreducible or connected fourth moment: $\langle 1234 \rangle = \langle 12 \rangle \langle 34 \rangle + \langle 13 \rangle \langle 24 \rangle + \langle 14 \rangle \langle 23 \rangle + \langle 1234 \rangle_c$.

The signal we are interested in, $\langle S_{\text{KSZ},i} S_{\text{KSZ},j} \rangle$, is directly related to the peculiar velocity logarithmic power spectrum Δ_v^2 . Here and below, $\Delta_Q^2 \equiv P_Q(k, z) k^3 / (2\pi^2)$, where k is the comoving wavenumber, z is the redshift and P_Q is the proper power spectrum of the random field Q . The variance is $\langle Q^2 \rangle = \int_0^\infty dk \Delta_Q^2 / k$.

The nonlinear scale at $z \sim 1$ is $k \simeq 0.5h/\text{Mpc}$, hence for wavenumbers $k < 1h/\text{Mpc}$ we can apply the weakly nonlinear perturbation theory. We also assume here that to first

order, the density and velocity are Gaussian random fields, related by the expression

$$\Delta_v(k, z) = f H a \Delta_m / H_0 k \sqrt{3}, \quad (6)$$

where a is the scale factor, $f \equiv d \ln D / d \ln a$, $H = H(z)$ is the Hubble parameter at redshift $z = 1/a - 1$ and $H_0 = H(0)$, and $D(z)$ is the usual linear growth factor for density perturbations. Δ_m^2 is the logarithmic power spectrum of the dark matter density fluctuations derived using the standard transfer function (Bardeen et al. 1986) and assuming $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $\sigma_8 = 0.9$ and $H_0 = 70 \text{ km/s/Mpc}$. Under these assumptions, to lowest order, the connected fourth moment vanishes because of Gaussianity while the nonlinear, mode coupling contribution to the signal is well approximated by the expression (Ma & Fry 2002)

$$\Delta_{\delta v}^2(\mathbf{k}) \approx \Delta_m^2(k, z) \sigma_v^2(z) / 2, \quad (7)$$

where $\sigma_v^2 = \int_0^\infty dk \Delta_v^2 / k$ is the line-of-sight velocity dispersion. We also assume that the two lines of sight are almost parallel. The accuracy of this assumption is better than $\sim 1\%$ for angular separations $< 10^\circ$.

Bulk flows are induced by the large scale gravitational potential, so the velocity field decouples from the small scale density field and the last expression in Eq.(7) remains valid and agrees with N-body simulations in the nonlinear regime, at redshifts $Z < 1$ and wavenumbers $k > 1h/\text{Mpc}$ if the linear Δ_m^2 is replaced by its nonlinear analog (Ma & Fry 2002).

To express these results in terms of the cluster number-density fluctuations $\delta_C(M)$, we will again turn to a simple Press-Schechter prescription, described earlier. The KSZ signal is then

$$\Delta_{\text{KSZ}}^2(k) = s_0^2 \left(\Delta_v^2(k) + \overline{b_n}^2 \Delta_{v\delta}^2 \right), \quad (8)$$

where

$$s_N \equiv \int_{M_*}^\infty b_n^N(M) S(M) (dn/dM) dM, \quad N = 0, 1,$$

and $\overline{b_n} \equiv s_1/s_0$. The quantities Δ_v^2 and $\Delta_{v\delta}^2$ are the logarithmic power spectra of spatial correlation functions $\langle v_1 v_2 \rangle$ and $\langle \delta_1 v_1 v_2 \delta_2 \rangle$, introduced above. To describe the density field, we use the cluster mass function, written as dn/dM , and M_* is the lower mass threshold of SZ selected clusters. For SPT, $M_* \sim 10^{14} M_\odot$. The redshift dependence of M_* (Holder et al. 2000) is weak and we assume a constant M_* . There is no biasing assumption for velocities because we assume that the galaxies trace the dark matter velocity field, in agreement with observations (Feldman, et al. 2003a). To calculate the effective cluster bias, we follow the halo model of (Mo & White 1996): $b_n(M) = 1 + (\nu^2 - 1)/(\delta_c/D)$, where $\delta_c \simeq 1.686$, $\nu = \delta_c/(D\sigma(M))$ and $D(z)$ is the linear density growth factor. The quantity $\sigma(M)$ is the rms density fluctuation in a spherical volume containing mass M and we have $\sigma(M_*) \simeq \sigma_8 = 0.9$. For the present estimate we use the approximation $\overline{b_n} = b_n(M_*)$. At present time $\overline{b_n}(M_*, 0) \approx 3$. This value is consistent with most recent measurements of the correlation function of optically selected galaxy clusters in the Sloan Digital Sky Survey (Estrada, Sefusati & Frieman 2008). This is, indeed, a simplified version of the biasing scheme, however for the scales of interest in this paper it is justified.

² <http://www.darkenergysurvey.org/>

³ The photo- z of each galaxy has dispersion ~ 0.05 . Clusters have $\gtrsim 100$ galaxies and thus the determined z dispersion is $\lesssim 0.005$.




Figure 1. The normalized KSZ cluster flux power spectra at $z = 0$ (upper black solid line) and $z = 1$ (lower blue solid line) respectively. The power spectrum, $\Delta_{\text{KSZ}}^2(k)$ is expressed in dimensionless units, defined in §2. The dashed and dotted lines show the contributions from the vv and $v\delta v\delta$ terms, respectively. The solid line shows the sum of these two terms. $\sigma_{v100}^2 \equiv \sigma_v^2/(100 \text{ km/s})^2$. The normalization is $\Delta_{\text{KSZ}}^2/s_0^2$.

In Fig. 1 we show the normalized KSZ flux power spectrum at $z = 0$ and at $z = 1$. It is clear that the vv contribution dominates at small k whereas the $\delta v\delta v$ term dominates in the high k tail. Δ_v^2 peaks at $k \sim 0.05 h/\text{Mpc}$, because of the k^2 denominator, while Δ_m^2 , associated with contaminants traced by point sources, keeps decreasing toward large scales. On the other hand, Δ_{KSZ}^2 increases with k because of the presence of the mode-coupling term. Note also, that compared to Δ_m^2 , Δ_{KSZ}^2 has a weak z dependence at $k \lesssim 0.03 h/\text{Mpc}$.

We have not considered the effect of redshift distortion. It reduces the power of the Δ_v^2 part by a factor of $\sim \exp[-k^2 \sigma_v^2/H^2(a)]$. Since $\sigma_v \sim 200 \text{ km/s}$, the redshift distortion can be neglected at $k \lesssim 0.2 h/\text{Mpc}$. On the other hand, the redshift distortion increases the power of the $\Delta_{v\delta}^2$ part in the linear regime (Kaiser 1987). Since $\Delta_{v\delta}^2$ dominates over Δ_v^2 at $k \gtrsim 0.1 h/\text{Mpc}$ (see Fig. 1), neglecting the redshift distortion tends to underestimate the signal Δ_{KSZ}^2 and thus overestimates the relative errors.

3 NOISE SOURCES

To study the detectability of Δ_{KSZ}^2 , let us consider the South Pole telescope experiment⁶, planned for the near future. The SPT will cover the 217 GHz range and about 4000 deg^2 with arc-minute resolution. It is expected to observe more than 20,000 clusters with masses greater than $10^{14} M_\odot$ for the thermal SZ signal. Virtually all the TSZ clusters have flux measurements at 220 GHz band, a fraction of this flux comes

from KSZ effects. The corresponding KSZ flux may be too weak to detect cluster by cluster, however, our method provides a way to extract this KSZ signal at 220 GHz band by statistically weighting all fluxes. Thus, cluster finding is done by combining all bands (90, 150, 220 and 270 GHz). Given these TSZ selected clusters, we can measure the CMB flux enclosed in the same region at 220 GHz. The measured flux is the sum of KSZ, residual TSZ, CMB and various other contaminations. Indeed, for many individual clusters, the KSZ signal is overwhelmed by other contaminants, however, the correlation approach described here is able to separate the KSZ signal from various sources of noise.

In the following we discuss the three dominant error sources in the following subsections. We assume redshift bin size $\Delta z = 0.2$ and k bin size $\Delta k = 0.4k$. The statistical errors scale as $\Delta k^{-1/2}$.

3.1 Diffuse foregrounds and backgrounds

Clusters have typical bulk velocity around several hundred kilometer per second, so the typical KSZ signal is $\Delta T_{\text{KSZ}} \sim 20 \mu\text{K} \langle \tau \rangle / 0.01$. Primary CMB, which is indistinguishable from the KSZ signal in frequency space, has intrinsic temperature fluctuation $\Delta T \sim 100 \mu\text{K}$. The cosmic infrared background (CIB) has mean $T \sim 20 \mu\text{K}$, if scaled to 217 GHz (Fixsen et al. 1998). A cluster KSZ filter can be applied to remove the mean backgrounds while keeping the KSZ signal. This filter must strongly match the cluster KSZ profile while having zero integrated area. Since the angular sizes of clusters are usually close to several arc-minutes, such a filter peaks at multipole l around several thousands and thus removes the dominant CMB signal, concentrated near $l \lesssim 1500$. For such filters, at $\sim 217 \text{ GHz}$, the galactic synchrotron, dust emission, free-free foregrounds, the radio and TSZ backgrounds are all negligible due to their frequency or scale dependence (see, e.g. (Wright 1998; ?)). Thus we only discuss the contaminations of the primary CMB, CIB and background KSZ.

The optimal filter can be constructed from the intracluster gas profile inferred from the TSZ survey. Without loss of generality, we choose the electron density profile as $n_e(r) \propto (1 + r^2/r_c^2)^{-1}$ and a matched filter $W_\ell = 6(\ell/\ell_f)^2 \exp(-(\ell/\ell_f)^2)$. For these particular choices, the filtered KSZ temperature, $\tilde{\Delta T}_{\text{KSZ}}$, peaks at $\ell_f \approx 1.1/\theta_c \approx 3800(1'/\theta_c)$, where θ_c is the angular core radius, and the peak value is $\tilde{\Delta T}_{\text{KSZ}} \simeq \Delta T_{\text{KSZ}} \simeq 9v_{100}[\langle \tau \rangle / 0.01] \mu\text{K}$. We adopt this ℓ_f and $\theta_c = 0.4 h^{-1} \text{ Mpc} / \chi(z)$ to estimate the noise. Here, $\chi(z)$ is the comoving angular distance.

The correlations of filtered backgrounds (with zero mean flux), originating from both their intrinsic and cluster correlations, are

$$\tilde{\xi}_b(r) \approx \left[\int \omega(M) dn(M) \right]^2 \left[1 + \overline{b_n}^2 \xi_m(r) \right] \tilde{w}_b(\theta), \quad (9)$$

where $\theta = r/\chi$, while $\tilde{w}_b(\theta)$ is the corresponding (filtered) background angular correlation function and $\omega(M)$ is the solid angle subtended by a cluster of mass M at redshift z . Finally, ξ_m is the dark matter correlation function.

The same cluster survey measures both $\int \omega dn$ and $\overline{b_n}^2 \xi_m$, while the CMB C_ℓ 's are known from observations (Hinshaw et al. 2003). Alternatively, for the cosmological model assumed here, the expected C_ℓ 's can be also calcu-




Figure 2. The dominant errors of the cluster KSZ power spectrum measurement. The solid lines are the errors caused by the cosmic variance and shot noise. The short dash lines are the systematic errors caused by the CIB and KSZ background. The long dash lines are the statistical errors caused by the primary CMB. The dot lines (which are visible only in the upper left panel) are the statistical errors caused by sources associated clusters. Systematic errors are sub-dominant at $k \lesssim 1 h/\text{Mpc}$.

lated from first principles (Seljak & Zaldarriaga 1996). Thus the CMB contamination can be easily subtracted from the correlation estimator, leaving the statistical errors from intrinsic CMB fluctuations over cluster regions as the only CMB contaminant.

In principle, the CIB and KSZ contaminations can be removed as well. But since both the amplitude and shape of CIB and KSZ power spectra are highly uncertain, we do not attempt to subtract their contribution from the correlation estimator. The CIB power spectrum is $C_\ell^{(1)} \ell^2 / (2\pi) \simeq (4\mu\text{K})^2 (\ell/10^3)^{0.7}$ and the KSZ power spectrum is $C_\ell^{(2)} \ell^2 / (2\pi) \simeq (2.7\mu\text{K})^2$ (Knox, Cooray & Eisenstein 2001). The resulting upper limit on the fractional systematic error is

$$\eta(k) \approx \frac{(1 + A_k) \sum C_\ell^{(J)} W_\ell^2 \ell^2}{2\pi B_k [9(\langle\tau\rangle/0.01)\mu\text{K}]^2}, \quad (10)$$

where $J = 1, 2$ is the summation index; the spherical harmonic number, $\ell = k\chi$, is fixed, $A_k \equiv \bar{b}_n^{-2} \Delta_m^2$, while $B_k \equiv \Delta_{v100}^2 + \bar{b}_n^2 \Delta_{v\delta100}^2$. Where $\Delta_{v\delta100}^2$ is the $\Delta_{v\delta}^2$ term, expressed in units of $(100 \text{ km/s})^4$. Clearly, $\eta(k)$ is at most several percent at $k \lesssim 1 h/\text{Mpc}$ (see Fig. 2).

3.2 Sources associated with clusters

The map filter does not remove the contaminations from sources associated with clusters. The flux of cluster radio and IR sources is $\sim 10^3 \text{ Jy/sr}$ at 217 GHz (Aghanim, Hansen & Lagache, 2005). Using multi-frequency information and

resolved source subtraction, one is likely able to subtract much of these contaminations. Rather conservatively, we assume that, at $\nu \sim 217 \text{ GHz}$, the total flux contributed by the sources associated with clusters is less than $\sim 5 \times 10^3 \text{ Jy/sr}$ at $z = 0.5$ and scale it to other redshifts assuming no intrinsic luminosity evolution. At 217 GHz, the non-relativistic TSZ vanishes. The relativistic correction of cluster TSZ effect shifts the cross over point slightly to higher frequency and thus in principle introduces a residual TSZ signal in $\sim 217 \text{ GHz}$ band. But the correction to ΔT is generally only several percent of the TSZ at Rayleigh-Jeans regime (Stebbins 1997; ?).

The mean flux of the cluster sources, S_{cl} , is subtracted in our estimator (Eq. 3). Since the cluster thermal energy, IR and radio flux should be mainly determined by local processes, one can omit the possible large-scale correlations of these quantities. Thus these sources do not cause systematic errors. But since the mean square of the flux fluctuations, caused by clusters, $\langle \delta S_{\text{cl}}^2 \rangle$, does not vanish, they do contribute a shot noise, which will be estimated in §3.3. Further, they contribute a statistical noise term proportional to the cluster clustering, whose expected fractional error is $\eta(k) \approx 2\pi A_k \langle \delta S_{\text{cl}}^2 \rangle / S_{100}^2 B_k \sqrt{V k^2 \Delta k}$, for a survey of volume V . This can be rewritten as

$$\eta(k) \approx 5 \times 10^{-3} \frac{\langle \delta S_{\text{cl}}^2 \rangle}{z S_{100}^2} \sqrt{\frac{\Delta z}{0.2} \frac{\Delta k/k}{0.4}}. \quad (11)$$

It is reasonable to assume that $\langle \delta S_{\text{cl}}^2 \rangle^{1/2} \lesssim \langle S_{\text{cl}} \rangle$. Thus the error caused by the sources associated with clusters is negligible at all scales and redshifts (See Fig. 2).

3.3 Cosmic variance and shot noise

The signal intrinsic cosmic variance dominates at large scales. The number of clusters is limited, so the shot noise is large, even in the linear regime. We believe that there are five dominant sources of shot noise. The shot noise power spectrum is then given by

$$\tilde{\Delta}_{\text{shot}}^2 = \frac{k^3}{2\bar{n}\pi^2} \sum_{J=1}^5 \tilde{\sigma}_J^2, \quad (12)$$

where the summation index J corresponds to mean square noise values of discrete sources, associated with clusters ($J = 1$), the instrumental noise, the residual flux fluctuations of primary CMB, CIB, and the background KSZ ($J = 2$ to 5, respectively). $\bar{n}$ is the mean number density of observed clusters, which can be calculated given the halo mass function, the survey specification and the gas model. We use the expression $\bar{n}(z) = 3 \times 10^{-5} / (1+z)^3 (h/\text{Mpc})^3$. We estimate the contributions of the five noise sources discussed here as follows: $\tilde{\sigma}_1 \approx 5\mu\text{K}$ and $\tilde{\sigma}_J \approx 20\mu\text{K}$ for $J = 2-5$. The filtered $\tilde{\sigma}$ of intracluster gas internal flow (Nagai, Kravtsov & Kosowsky 2003) is $\lesssim 10\mu\text{K}$, due to its random nature. Lack of knowledge on its spatial distribution, we neglect this term. For SPT, the errors caused by the cosmic variance and shot noise ($\sim 10\%$) dominate over all other errors. The systematic errors are virtually always sub-dominant. In this sense, our method has the special advantage to measure the cluster peculiar velocity. For a future all sky survey, total error can be reduced to several percent level.

4 CONCLUSIONS

We presented a new method to estimate the peculiar velocity power spectrum using the KSZ effect. The signal (cluster bulk velocity correlation) is amplified by direct auto correlation of the cluster KSZ flux. Consequently, many systematics, such as internal flow involved in the usual v_p inversion, vanish and the majority of remaining systematics can be simply subtracted. The correlation method has the special advantage that statistical error dominates over systematics at effectively all k and z range.

Ideas similar to those expressed here about prospects of probing the peculiar velocity field of clusters at high redshift through KSZ flux correlations have been recently presented in (Hernandez-Monteagudo et al. 2006). Our papers are complementary: we focus on spatial correlations, while (Hernandez-Monteagudo et al. 2006) study angular correlations.

Our calculations show that the SPT can measure Δ_v^2 to $\sim 10\%$ accuracy at $k \sim 0.02\text{--}0.3h/\text{Mpc}$ and a future all sky survey can improve this measurement by a factor of few. These results should be regarded as preliminary – the goal of this paper was to present a new idea and a rough estimate of the expected signal-to-noise ratio. A comparison with more realistic numerical simulations should be the next step.

5 ACKNOWLEDGMENTS

PJZ thanks the support of the DOE and the NASA grant NAG 5–10842 at Fermilab at the early stage of this work and the National Science Foundation of China grants 10673022 and 10533030. HAF was supported in part by the University of Kansas GRF and by a grant from the Research Corporation. RJ was supported by (to be added just before submission). AS is supported by the DOE and the NASA grant NAG 5–10842 at Fermilab.

6 REFERENCES

REFERENCES

Aghanim, N., Gorski, K.M., & Puget, J. L., 2001, *A&A*, 374, 1
Aghanim, N., Hansen, S., & Lagache, G., 2005, *A&A*, 439, 901A
Atrio-Barandela, F., Kashlinsky, A., & Mucket, J.P., 2004, *ApJ*, 601, L111
Bardeen, J.M., Bond, J. R., Kaiser, N. & Szalay, A. S., 1986, *ApJ*, 304, 15
Bennett, C.L., et al. 2003, *ApJs*, 148, 97
Bhattacharya, S. and Kosowsky, A., 2007, *ApJ*, 659, L83
Bonvin, C., Durrer, R., & Kunz, M. 2006, *PRL*, 96, 191302
Courteau, S., Willick, J., Eds., 2000, *Cosmic Flows 1999*, ASP Conference Series Vol. 201, Sheridan Books, Astron. Soc. Pac., San Francisco.
da Costa, L. N., Nasser, A., Freudling, W., Giovanelli, R., Haynes, M. P., Salzer, J. J., Wegner, G., 1998, *MNRAS*, 299, 425
Diaferio, A., et al. 2005, *MNRAS*, 356, 1477
Estrada, J., Sefusati, E. & Frieman, J., 2008, *arXiv:0801.3485* [astro-ph]
Feldman, H., et al. 2003a, *ApJ*, 596, L131
Feldman, H. A., Watkins, R. Melott, A. L. & Chambers, S. W., 2003b, *ApJ*, 599, 820
Feldman, H. A. & Watkins, R., 2008, *MNRAS*, 387, 825

Fixsen, D. J., Dwek, E., Mather, J. C., Bennett, C. L., Shafer, R. A., 1998, *ApJ*, 508, 123
Fosalba, P. & Doré, O., 2007 *PRD*, 76 103523
Gordon, C., Land, K. and Slosar, A., 2007, *PRL*, 99, 081301
Groth, E.J., Juskiewicz, R., & Ostriker, J.P., 1989, *ApJ*, 346, 558
Haehnelt, M.G., & Tegmark, M., 1996, *MNRAS*, 279, 545
Haugboelle, T., Hannestad, S., Thomsen, B., Fynbo, J., Sollerman, J., & Jha, S., 2007, *ApJ*, 661, 650
Hernandez-Monteagudo, C., Verde, L., Jimenez, R. & Spergel, D. N. 2006, *ApJ*, 643, 598
Hinshaw, G., et al. 2003, *ApJS*, 148, 135
Holder, G. P., Mohr, J. J., Carlstrom, J. E., Evrard, A. E., Leitch, E. M., 2000, *ApJ*, 544, 629
Holder, G., 2004, *ApJ*, 602, 18
Hui, L., & Greene, P. B., 2006, *PRD*, 73, 123526
Itoh, N., Kohyama, Y., & Nozawa, S. 1998, *ApJ*, 502, 7
Kaiser, N., 1987, *MNRAS*, 227, 1
Kashlinsky, A., 1987, *ApJ*, 317, 19
Kashlinsky, A., 1991, *ApJ*, 376, L5
Kashlinsky, A., & Atrio-Barandela, F., 2000, *ApJ*, 536, L67
Knox, L., Cooray, A. & Eisenstein, D., 2001, *ApJ*, 550, 7
Knox, L., Holder, G., & Church, S., 2004, *ApJ*, 612, 96
Ma, C.-P., & Fry, J.N., 2002, *Phys. Rev. Lett.*, 88, 211301
Mo, H. J. & White, S. D. M. 1996, *MNRAS*, 282, 347
Nagai, D., Kravtsov, A.V., & Kosowsky, A. 2003, *ApJ*, 587, 524
Neill, J. D., Hudson, M. J., & Conley, A. 2007, 661, *ApJL* 123
Ruhl, Z. et al., 2004, *Proc. SPIE*, Vol. 5498, p 11-29
Sarkar, D., Feldman, H.A. & Watkins, R., 2007, *MNRAS*, **375**, 69
Seljak, U., & Zaldarriaga, M., 1996, *ApJ*, 469, 437
Stebbins, A., 1997, *astro-ph/9705178*
Strauss, M.A., & Willick, J.A., 1995, *Phys. Rep.* 261, 271
Sunyaev, R. A. & Zeldovich, I. B., 1980, *MNRAS*, 190, 413
Vittorio, N., Juskiewicz, R., & Davis, M., 1986, *Nature*, 323, 132
Wang, L.F. 2007, *arXiv:0705.0368* [astro-ph]
Watkins, R. & Feldman, H.A. 2007, *MNRAS*, 379, 343-348
Wright, E., 1998, *ApJ*, 496, 1
Yoshida, N., Sheth, R. & Diaferio, A. 2001, *MNRAS*, 328, 669
Zel'dovich, Y.B. & Sunyaev, R., 1969, *Ap&SS*, 4, 301
Zhang, P., Stebbins, A., Juskiewicz, R. & Feldman, H. A., 2004 *arXiv:astro-ph/0410637*